

HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Medicine

journal homepage: <http://ees.elsevier.com/apjtm>Original research <http://dx.doi.org/10.1016/j.apjtm.2015.10.012>***Vibrio* spp. from *Macrobrachium amazonicum* prawn farming are inhibited by *Moringa oleifera* extracts**

Raimunda Sâmia Nogueira Brilhante^{1*}, Jamille Alencar Sales², Celia Maria de Souza Sampaio², Francisco Geraldo Barbosa³, Manoel de Araújo Neto Paiva², Glaucia Morgana de Melo Guedes¹, Lucas Pereira de Alencar², Yago Brito de Ponte², Tereza de Jesus Pinheiro Gomes Bandeira¹, José Luciano Bezerra Moreira¹, Débora de Souza Collares Maia Castelo-Branco¹, Waldemiro de Aquino Pereira-Neto¹, Rossana de Aguiar Cordeiro¹, José Júlio Costa Sidrim¹, Marcos Fábio Gadelha Rocha^{1,2}

¹Department of Pathology and Legal Medicine, Postgraduate Program in Medical Microbiology, Specialized Medical Mycology Center, Federal University of Ceará, Fortaleza, Ceará, Brazil

²School of Veterinary Medicine, Postgraduate Program in Veterinary Sciences, State University of Ceará, Fortaleza, Ceará, Brazil

³Department of Organic and Inorganic Chemistry, Federal University of Ceará, Fortaleza, Ceará, Brazil

ARTICLE INFO

Article history:

Received 15 Aug 2015

Received in revised form 20 Sep 2015

Accepted 30 Sep 2015

Available online 9 Oct 2015

Keywords:

Prawn

*Macrobrachium amazonicum**Moringa oleifera**Vibrio*

Antimicrobial

ABSTRACT

Objective: To investigate the *in vitro* antimicrobial potential of extracts of stem, leaves, flowers, pods and seeds of *Moringa oleifera* (*M. oleifera*) against *Vibrio* spp. from hatchery water and the prawn *Macrobrachium amazonicum*.

Methods: The ethanol extracts of stem, leaves, pods and seeds and chloroform extract of flowers of *M. oleifera* were tested against *Vibrio cholerae* (*V. cholerae*) serogroups non-O1/non-O139 ($n = 4$), *Vibrio vulnificus* ($n = 1$) and *Vibrio mimicus* ($n = 1$). *Escherichia coli* (*E. coli*) (ATCC[®] 25922) was used as quality control. *Vibrio* species were obtained from *Macrobrachium amazonicum* prawns and from hatchery water from prawn farming. The Minimum Inhibitory Concentration (MIC) was determined by broth microdilution method.

Results: The best result was obtained with the ethanol extract of pods, which inhibited three strains of the *V. cholerae*, *Vibrio vulnificus*, *Vibrio mimicus* and *E. coli* (MIC range 0.312–5.000 mg/mL). The chloroform extract of flowers was effective against all *V. cholerae* strains and *E. coli* (MIC range 0.625–1.250 mg/mL). However, the ethanol extracts of stem and seeds showed low effectiveness in inhibiting the bacterial growth.

Conclusions: The extracts of pods, flowers and leaves of *M. oleifera* have potential for the control of *Vibrio* spp. Further studies are necessary to isolate the bioactive compounds responsible for this antimicrobial activity.

1. Introduction

The cultivation of shrimp can be threatened by diseases caused by *Vibrio* species, which can result in up to 100% mortality, 24 h after the appearance of infection [1]. *Vibrio cholerae* (*V. cholerae*), *Vibrio mimicus* (*V. mimicus*) and *Vibrio vulnificus* (*V. vulnificus*) are opportunistic pathogens

capable of causing lethal infections in farmed crustaceans when there are stressful environmental conditions, nutritional imbalance and predisposing lesions [2]. Moreover, antimicrobial resistance in these microorganisms has been observed [3].

The emergence of antibiotic resistant bacteria has driven research to find new compounds with antimicrobial properties in plants used in traditional medicine, such as *Moringa oleifera* (*M. oleifera*) (Lam.) [4–10].

M. oleifera is a well-known and widely distributed tree species, belonging to the family Moringaceae [11]. In Brazil it can be found in the Northeast, mainly in the states of Maranhão, Piauí and Ceará [9]. The antimicrobial properties of *M. oleifera* have been attributed to different parts of the plant, such as leaves, flowers, seeds, pods and stems [9,12,13]. The

*Corresponding author: R.S.N. Brilhante. Rua Coronel Nunes Melo, s/n, Rodolfo Teófilo. CEP: 60.430-270. Fortaleza, CE, Brazil.

Tel: +55 (85) 3366 8319

E-mail: brilhante@ufc.br

Peer review under responsibility of Hainan Medical College.

Foundation project: This work was supported by grants from the National Council for Scientific and Technological Development (CNPq; Brazil; Processes 445670/2014-2) and Coordination Office for the Improvement of Higher Education Personnel (AEI-0052-000650100/11).

literature reports the antimicrobial potential of *Moringa* against bacteria and fungi isolated from shrimp farming [5,9].

Thus, the objective of this study was to evaluate the *in vitro* antimicrobial potential of extracts of stem, leaves, flowers, pods and seeds of *M. oleifera* against *Vibrio* species isolated from hatchery water and *Macrobrachium amazonicum* (*M. amazonicum*) prawn.

2. Materials and methods

2.1. Extracts

The extracts were obtained from specimens of *M. oleifera* grown in Fortaleza, Ceará, Brazil, and provided by the Laboratory of Applied Phytochemistry, Federal University of Ceará. Stem, leaves, pods and seeds were dried in a heated chamber at 40 °C and then subjected to three successive extractions by cold maceration with ethanol at intervals of 24 h, originating the ethanol extracts, while flowers were dried at 40 °C and then subjected to three successive extractions by cold maceration with chloroform at intervals of 24 h, originating the chloroform extract. After filtration, the respective solvents were evaporated under reduced pressure in a rotary evaporator, leaving only the concentrated constituents extracted from the plant parts [7].

2.2. Strains of *Vibrio* spp.

V. cholerae serogroups non-O1/non-O139 ($n = 4$), *V. mimicus* ($n = 1$) and *V. vulnificus* ($n = 1$), belonging to the bacterial collection of the Laboratory of Emerging and Re-emerging Pathogens of Ceará Federal University, were used in this study. These strains were obtained through the collection of specimens of ovigerous *M. amazonicum* females from Sapiranga Lake (3°48'3.46" S and 38°27'30.83" W), Fortaleza, Ceará, Brazil, and samples of hatchery water from *M. amazonicum* farming, during the larval development stage, at the Laboratory of Shrimp Farming of the State University of Ceará.

2.3. *In vitro* susceptibility test

The *in vitro* susceptibility test with extracts of *M. oleifera* was performed following the method described by Rocha [7] in 2011, with some modifications. Initially, each extract was dissolved in dimethyl sulfoxide (DMSO) (LGC Biotecnologia Ltda, São Paulo, Brazil) and then diluted in Müeller-Hinton broth (Difco™, São Paulo, Brazil). In previous tests with strains of *Vibrio* spp., we verified that DMSO alone at concentrations up to 5% was not able to inhibit the growth of these strains. Thus, the concentration of DMSO in the

susceptibility test did not exceed 5%, to assure that results refer to the action of each extract tested [7].

Minimum inhibitory concentration (MIC) of *M. oleifera* extracts against strains of *Vibrio* spp. was determined through the broth microdilution method as standardized by the Clinical Laboratory Standards Institute, based on the document M07-A9 [14]. The MIC was considered the lowest concentration of the extracts able to inhibit 50% of bacterial growth compared to the control growth [9]. The strain *Escherichia coli* (*E. coli*) (ATCC® 25922) was included as quality control. The initial concentration of each extract used was 20 mg/mL and the range of concentrations evaluated in the susceptibility test was from 0.01 to 5.00 mg/mL. The microdilution assays were performed in 96 well plates with a final volume of 200 µL, incubated at 35 °C and read after 20 h, according to the document M45-A2 [15]. The inocula were prepared at a turbidity of 0.5 on McFarland scale (108 CFU/mL), and then diluted with Müller-Hinton broth so that each well after inoculation presented approximately 5×10^5 CFU/mL. All assays were performed in duplicate, and for each strain growth control and sterility control of the culture medium were included [14]. The reading was performed with a spectrophotometer (BioteK®, Winooski, United State) at 590 nm and the obtained absorbance values were corrected by the absorbance obtained for each tested extract alone. Only extracts that inhibited the control strain (*E. coli* ATCC® 25922) were considered to have antimicrobial activity against *Vibrio* spp. Chloramphenicol (Sigma-Aldrich® Brazil Ltda, São Paulo, Brazil) was used as standard antibiotic, as recommended by the document M100-S22 [16].

2.4. Research licensing

This study was previously approved by the Chico Mendes Institute for Conservation of Biodiversity/Biodiversity Authorization and Information System – SISBIO, under the number 28175-1.

3. Results

The ethanol pod extract showed the best inhibitory activity against isolates from the hatchery water, with MIC values ranging from 0.3125 to 1.250 mg/mL against three strains of *V. cholerae* non-O1/non-O139 (3/4) and an MIC of 5 mg/mL against *V. vulnificus*. This extract also showed inhibitory effect against *V. mimicus* from the digestive tract of *M. amazonicum* and *E. coli* (ATCC® 25922), with MIC values of 1.25 mg/mL and 2.50 mg/mL, respectively (Table 1).

Table 1

MIC values of *Moringa oleifera* extracts against *Vibrio* spp. recovered from hatchery water and *Macrobrachium amazonicum* prawn.

Source	Species	MIC (mg/mL)					MIC (µg/mL)
		Ethanol stem	Ethanol leaves	Chloroform flowers	Ethanol pods	Ethanol seeds	Chloramphenicol
Hatchery water	<i>V. cholerae</i>	n.i	0.078	1.25	0.312 5	5	0.5
	<i>V. cholerae</i>	n.i	0.625	0.625	n.i	n.i	0.5
	<i>V. cholerae</i>	n.i	n.i	0.625	1.25	n.i	0.5
	<i>V. cholerae</i>	2.5	n.i	0.625	0.312 5	2.5	0.5
	<i>V. vulnificus</i>	n.i	n.i	n.i	5	n.i	0.5
Prawn	<i>V. mimicus</i>	1.25	5	n.i	1.25	n.i	0.5
	<i>E. coli</i> ATCC®M25922	n.i	5	1.25	2.5	n.i	4

The chloroform extract of flowers was effective against all strains of *V. cholerae* non-O1/non-O139, with MIC values ranging from 0.625 to 1.250 mg/mL, as well as against *E. coli* strain (ATCC® 25922) (MIC 1.25 mg/mL) (Table 1).

The ethanol extract of leaves, in turn, had inhibitory activity against two strains of *V. cholerae* non-O1/non-O139 (2/4), with MIC values of 0.078 and 0.625 mg/mL, and also presented MIC value of 5 mg/mL against *V. mimicus* and *E. coli* (Table 1).

The ethanol extract of stem and seeds showed low effectiveness in inhibiting the growth of *Vibrio* spp. strains, as well as no effect against *E. coli*, as shown in Table 1. The assay performed with the standard antibiotic, chloramphenicol, against the control strain *E. coli* (ATCC® 25922) presented MIC value within the range established in the document M100-S22 (17) (Table 1).

4. Discussion

Almost all parts of the *M. oleifera* have multiple industrial and medical uses [17]. The pharmacological potential of this plant has been described in the literature, in particular the antimicrobial activity of the extracts of pods, flowers, leaves, stem and seeds [9,10,18]. Thus, this study aimed to verify the antimicrobial activity of *M. oleifera* against *Vibrio* species, considering that these microorganisms are opportunistic zoonotic pathogens and can cause economical losses to shrimp farming and public health problems, as well [19].

The results of this study demonstrate the antibacterial activity of extracts from different parts of *M. oleifera* against *Vibrio* strains isolated from prawn hatchery water and from *M. amazonicum*. The antibacterial activity of the extracts of leaves, flowers and pods has been reported in other studies against Gram-negative bacteria [10,11]. In addition, the antimicrobial activity of *M. oleifera* extracts has also been observed against *Candida* species isolated from *M. amazonicum* farming in previous study [9].

The extracts of pods, flowers and leaves of *M. oleifera* showed to be more effective, when compared to the other extracts. The ethanol extract of pods presented the best inhibitory activity against species of *Vibrio* spp., and it was the only one capable of inhibiting *V. vulnificus*. The growth of the *E. coli* strain (ATCC® 25922) was also inhibited by this extract. Arora and Onsare [10] verified the effectiveness of the pod extract of *M. oleifera* against the Gram-negative bacteria *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *E. coli* and *Salmonella typhimurium*. However, there are few reports regarding the inhibitory effect of extracts of *M. oleifera* against *Vibrio* species.

Some studies on the history of traditional medicine demonstrated the effectiveness of preparations made of leaves, pods, flowers and seeds against microorganisms that cause human infections [17]. Therefore, studies with *V. mimicus*, *V. vulnificus* and *V. cholerae* non-cholera serogroup are important, because these bacteria inhabit estuarine and marine environments and have often been associated with sporadic cases of diarrhea, sepsis and infections, after ingestion of seafood or exposure to contaminated aquatic environment [19–21].

In a previous study by Rocha [9] in 2014, the chloroform extract of flowers presented antifungal activity against *Candida* spp. and *Hortaea werneckii* isolated from *M. amazonicum* farming. Based on the observation of this antimicrobial action, we tested the chloroform extract of flowers against *Vibrio* species from the same environment.

The chloroform extract of flowers also inhibit *Vibrio* strains from *M. amazonicum* cultivation water and the control strain. According to Anwar [11], the antibacterial property of *M. oleifera* flowers has been attributed to a substance called pterygospermin [22]. In addition to this compound, other substances, such as benzyl glucosinolates and their cognate isothiocyanate extracted from *M. oleifera* also have antimicrobial properties against bacteria [17]. Therefore, it is worth noting the need to investigate the bioactive compounds of plants that exhibit antimicrobial activity.

The low antimicrobial effectiveness of extracts of seeds and stem of *M. oleifera* found in this study has also been demonstrated by other authors [9,13,23]. Probably, this low antimicrobial effectiveness can be associated with inherent factors of *Vibrio* spp. which decrease or neutralize the activity of the bioactive compounds contained in the seeds and stem of *M. oleifera* [24].

Thus, the ethanol extracts of pods and leaves and the chloroform extract of flowers of *M. oleifera* presented potential to control *Vibrio* spp., although further research is necessary to determine the bioactive compounds responsible for this antimicrobial activity.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgments

This work was supported by grants from the National Council for Scientific and Technological Development (CNPq; Brazil; Processes 445670/2014-2) and Coordination Office for the Improvement of Higher Education Personnel (AEI-0052-000650100/11).

References

- [1] Costa RA, Vieira GHF, Silva GC, Vieira RHSF, Sampaio SS. Susceptibilidade 'in vitro' a antimicrobianos de estirpes de *Vibrio* spp isoladas de camarões (*Litopenaeus vannamei*) e de água de criação destes animais provenientes de uma fazenda de camarões no Ceará. *Braz J Vet Res An Sci* 2008; **45**(6): 458-462.
- [2] Banerjee S, Ooi MC, Shariff M, Khatoon H. Antibiotic resistant *Salmonella* and *Vibrio* associated with farmed *Litopenaeus Vannamei*. *Sci World J* 2012; **2012**(1): 130-136.
- [3] Reboças RH, Sousa OV, Lima AS, Vasconcelos FR, Carvalho PB, Vieira RHSF. Antimicrobial resistance profile of *Vibrio* species isolated from marine shrimp farming environments (*Litopenaeus vannamei*) at Ceará, Brazil. *Environ Res* 2011; **111**(1): 21-24.
- [4] Chuang PH, Lee CW, Chou JY, Murugan M, Shieh BJ, Chen HM. Anti-fungal activity of crude extracts and essential oil of *Moringa oleifera* Lam. *Bioresour Technol* 2007; **98**(1): 232-236.
- [5] Vieira GHF, Mourão JA, Ângelo AM, Costa RA, Vieira RHSF. Antibacterial effect (in vitro) of *Moringa oleifera* and *Annona muricata* against Gram positive and Gram negative bacteria. *J Inst Trop Med São Paulo* 2010; **52**(3): 129-132.
- [6] Arun T, Rao PCH. Phytochemical screening and antibacterial activity of *Moringa oleifera* Lam. against *Proteus mirabilis* from urinary tract infected patients. *Int J PharmTech Res* 2011; **3**(4): 2118-2123.
- [7] Rocha MFG, Aguiar FLN, Brilhante RSN, Cordeiro RS, Teixeira CEC, Castelo-Branco DSCM, et al. Extratos de *Moringa oleifera* e *Vernonia* sp. sobre *Candida albicans* e *Microsporium canis* isolados de cães e gatos e análise da toxicidade em *Artemia* sp. *Cienc Rural* 2011; **41**(10): 1807-1812.

- [8] Prasad TNVKV, Elumalai EK. Biofabrication of Ag nanoparticles using *Moringa oleifera* leaf extract and their antimicrobial activity. *Asian Pac J Trop Biomed* 2011; **1**(6): 439-442.
- [9] Rocha MFG, Alencar LP, Brilhante RSN, Sales JA, Ponte YB, Rodrigues PHA, et al. *Moringa oleifera* inhibits growth of *Candida* spp. and *Hortaea werneckii* isolated from *Macrobrachium amazonicum* prawn farming with a wide margin of safety. *Cienc Rural* 2014; **44**(12): 1-7.
- [10] Arora DS, Onsare JG. *In vitro* antimicrobial evaluation and phytoconstituents of *Moringa oleifera* pod husks. *Ind Crops Prod* 2014; **52**(1): 125-135.
- [11] Anwar F, Latif S, Ashraf M, Gilani AH. *Moringa oleifera*: a food plant with multiple medicinal uses. *Phytother Res* 2007; **21**(1): 17-25.
- [12] Ferreira RS, Napoleão TH, Santos AFS, Sá RA, Carneira-da-Cunha MG, Morais MMC, et al. Coagulant and antibacterial activities of the water-soluble seed lectin from *Moringa oleifera*. *Appl Microbiol* 2011; **53**(2): 186-192.
- [13] Arora DS, Onsare JM, Kuar H. Bioprospecting of *Moringa* (Moringaceae): microbiological perspective. *J Pharmacogn Phytochem* 2013; **1**(6): 193-215.
- [14] Clinical and Laboratory Standards Institute. *Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically*. CLSI document M7-A9. 9 th ed. Wayne, PA: Clinical and Laboratory Standards Institute; 2012.
- [15] Clinical and Laboratory Standards Institute. *Methods for antimicrobial dilution and disk susceptibility testing of infrequently isolated or fastidious bacteria*. CSLI document M45-A2. 2 nd ed. Wayne, PA: Clinical and Laboratory Standards Institute; 2010.
- [16] Clinical and Laboratory Standards Institute. *Performance standards for antimicrobial susceptibility testing: Twenty – second information supplement*. CLSI document M100-S22. Wayne, PA: Clinical and Laboratory Standards Institute; 2012.
- [17] Fahey JW. *Moringa oleifera*: a review of the medical evidence for its nutritional, therapeutic, and prophylactic properties part 1. *Trees Life J* 2005; **1**(5): 1-15.
- [18] Onsare JG, Kaur H, Arora DS. Antimicrobial activity of *Moringa oleifera* from different locations against some human pathogens. *Acad J Med Plant* 2013; **1**(5): 80-91.
- [19] Austin B. Vibrios as casual agentes of zoonoses. *Vet Microbiol* 2010; **140**(3-4): 310-317.
- [20] Igbiosa EO, Okoh AI. Emerging *Vibrio* species: an unending threat to public health in developing countries. *Res Microbiol* 2008; **159**(7-8): 495-506.
- [21] Ottaviani D, Leoni F, Rocchegiani E, Santarelli S, Masini L, Di Trani V, et al. Prevalence and virulence properties of non-O1, non-O139 *Vibrio cholerae* strains from seafood and clinical samples collected in Italy. *Int J Food Microbiol* 2009; **132**(1): 47-53.
- [22] Das BR, Kurup PA, Rao Narasimha PL. Antibiotic principle from *Moringa pterygosperma*. VII. Antibacterial activity and chemical structure of compounds related to pterygospermin. *Indian J Med Res* 1957; **45**(2): 191-196.
- [23] Jabeen R, Shahid M, Jamil A, Ashraf M. Microscopic evaluation of the antimicrobial activity of seed extracts of *Moringa oleifera*. *Pak J Bot* 2008; **40**(4): 1349-1358.
- [24] Singh RSG, Negi PS, Radha C. Phenolic composition, antioxidant and antimicrobial activities of free and bound phenolic extracts of *Moringa oleifera* seed flour. *J Funct Foods* 2013; **5**(4): 1883-1891.